

Full length Research paper

Impacts of dietary tyrosine on serum cholesterol portions in rodents

Chris DW¹, Messy B² and Adams N^{3*}

¹Department of Biochemistry, University of Tennessee Institute of Agriculture, Smith International Center, 2640 Morgan Circle Drive, Knoxville, TN 37996 USA

²Zamura Feeds, Inc., Ruhungeri, Musanze District, Rwanda

Accepted 22 March, 2019

The current investigation was attempted to gauge the impacts of dietary tyrosine included to rodent diet serum cholesterol levels in the rodent. A sum of twenty Wistar strain pale skinned person rodents were taken care of with various portions of tyrosine improved eating regimens containing 0.8 g/100 g, 1.0 g/100 g and 1.2 g/100 g. Following 3 weeks of trial taking care of, there was huge increment ($p < 0.05$) altogether postprandial serum cholesterol of rodents took care of with evaluated of tyrosine when contrasted and the ordinary control. Same pattern was followed in the week 2 of a similar taking care of example. The impacts of dietary tyrosine supplementation on cholesterol levels of the high thickness lipoprotein (HDL) portion were practically identical, yet not all noteworthy on the week 3 treatment. In any case, there was noteworthy reduction ($p < 0.05$) in week 2 of rodents took care of with the diverse reviewed portions of the tyrosine dinner when contrasted and the ordinary benchmark group. Moreover, noteworthy increment was likewise seen in the low thickness lipoprotein (LDL) when contrasted with the control after week 2 and 3 of tyrosine dinner treatment. These outcomes uncovered that tyrosine supplementation in a physiological sum may expand cholesterol levels in the rodent when added to consume less calories, with a moderate arrival of tyrosine during absorption.

Key words: Impacts Dietary tyrosine, Postprandial serum cholesterol, high density lipoprotein (HDL), low density lipoprotein (LDL), Rodents.

INTRODUCTION

Food helps humans maintain good health by providing all essential nutrients. Consuming a variety of foods in balanced proportions will prevent deficiency diseases and chronic diet-related disorders. Amino acids have many functions in the body. They are the building

blocks for all body protein - structural proteins that build muscle, connective tissues, bones and other structures, and functional proteins in the form of thousands of metabolically active enzymes (Elwes et al., 1989; Fernstrom, 2000). Amino acids provide the body with the nitrogen that is essential for growth and maintenance of all tissues and structures. Proteins and amino acids also serve as a source of energy, providing about 4 calories per gram. Aside from these general

Corresponding author's Email: adam123@gmail.com

Table 1: Composition of basal diet (Growers feed, Grand Cereals LTD, Enugu).

Diet composition	Amount (%)
Crude protein	15
Fat	7
Crude fibre	19
Calcium	1.0
Phosphorus	0.35
Metabolisable energy	2550 kcal/kg

functions, individual amino acids also have specific functions in many aspects of human physiology and biochemistry (Salter, 1989).

Tyrosine is a non essential amino acid the body makes from another amino acid called phenylalanine. It is a building block for several important brain chemicals called neurotransmitters, including epinephrine, norepinephrine and dopamine (Moller et al., 1995). Neurotransmitters help nerve cells communicate and influence mood (Thomas et al., 1999). Tyrosine also helps produce melanin, the pigment responsible for hair and skin colour. It helps in the function of organs responsible for making and regulating hormones, including the adrenal, thyroid and pituitary glands (Shurtleff et al., 1994). It is involved in the structure of almost every protein in the body. It is rare to be deficient in tyrosine. Low levels have been associated with low blood pressure, low body temperature and an underactive thyroid (Sole et al., 1985; Deijen and Orlebeke, 1994; Deijen et al., 1999).

Cholesterol is insoluble in the blood; it must be attached to certain protein complexes called lipoproteins in order to be transported through the bloodstream. Low-density lipoproteins (LDLs) transport cholesterol from its site of synthesis in the liver to the various tissues and body cells, where it is separated from the lipoprotein and is used by the cell (Gordon et al., 1989). Cholesterol attached to LDLs is primarily that which builds up in atherosclerotic deposits in the blood vessels hence LDLs are termed 'bad' cholesterol (Olson, 1998). High-density lipoproteins (HDLs) may possibly transport excess or unused cholesterol from the tissues back to the liver, where it is broken down to bile acids and is then excreted thereby serving to retard or reduce atherosclerotic buildup, thus, it is termed 'good' cholesterol (Lewis and Rader, 2005).

When we take a close look at the diet of depressed people, an interesting observation is that their nutrition is far from adequate. They make poor food choices and select foods that might actually contribute to depression. Salter (1989) has reported that dietary tyrosine aids to reduce stress among troops. Recent evidence by Tumilty et al. (2011) suggests that oral tyrosine supplementation improves exercise capacity in athletes. A lot of research

has been carried out on tyrosine. This study is aimed at determining the effects of dietary tyrosine added to rat diet on serum cholesterol levels in rats.

MATERIALS AND METHODS

Reagents

L-tyrosine used in this study was sourced from Sigma Aldrich USA (Lot# SI bb7526V). All other chemicals and reagents used are of analytical grade.

Animals

A total of twenty (20) Wistar strain albino rats weighing between 134 -180 g bred in the animal house of the Department of Zoology, University of Nigeria Nsukka, were used in the experiment. The animals were kept under room temperature and were acclimatized in the new environment for a period of 7 days and fed non purified diet with the following diet composition as shown in Table 1 before the addition of the dietary tyrosine. The use of animal for research studies was ethically approved by the authorized committee of animal ethics, Department of Biochemistry, University of Nigeria, Nsukka.

Experimental design

After the acclimatization period, a total of 20 rats were used for the experiment and was divided into four groups consisting of four rats in each group as follows: Group 1: Control were fed basal diet; Group 2: Rat fed with 0.8 g/100 g of tyrosine diet; Group 3: Rat fed with 1 g/100 g of tyrosine diet; Group 4: Rat fed with 1.2 g/100 g of tyrosine diet.

The treatment lasted for twenty one (21) days in which blood samples of the rats were analysed on day 0, 14, and 21. The animals were anaesthetised and blood sample were collected through ocular puncture for biochemical analysis. Blood samples were received into clean dry centrifuge tube and left to clot at room temperature, then centrifuged at 33.5 g for 15 min to obtain the serum. The serum was carefully separated into dry clean Wassermann tubes, using a Pasteur pipette and kept frozen at (-20°C) until estimation of some biochemical parameters.

Cholesterol determination

Total cholesterol was determined according the method of Abell et al. (1952) as outlined in commercially available kits. Determination of the concentration of the serum total HDL and LDL was determined as described by Kameswara et al. (1999).

Cholesterol determination

The method of Abell et al. (1952) was followed.

Principle

Cholesterol was determined after enzymatic hydrolysis and oxidation. The indicator quinoneimine is formed from hydrogen peroxidase and 4-aminoantipyrine in the presence of phenol and peroxidase.

Test procedure

Three (3) test tubes were set up in a test tube rack and labeled blank, standard and sample, respectively. To the blank, was added (10 μ l) distilled H₂O, 10 μ l standard specimen was added to the standard test tube and 10 μ l sample (serum) was added to the sample test tube. To each of these test tubes was added 1000 μ l of the cholesterol reagent. It was thoroughly mixed and incubated for 10 min at room temperature (20-25°C). The absorbance of the sample (A_{sample}) against the blank was taken within 60 min at 500 nm.

Low density lipoprotein (LDL)

Principle

LDL-C can be determined as the difference between total cholesterol and the cholesterol content of the supernatant after precipitation of the LDL fraction by polyvinyl sulphate (PVS) in the presence of polyethyleneglycol monomethyl ether.

Procedure

The serum samples were kept at 2-8°C. The precipitant solution (0.1 ml) was added to 0.2 ml of the serum sample and mixed thoroughly and allowed to stand for 15 min. This was centrifuged at 2,000 xg for 15 min. The cholesterol concentration in the supernatant was determined. The concentration of the serum total cholesterol as described by Kameswara et al. (1999) was used.

Calculation

LDL-C (mmol/L) = Total cholesterol (mmol/L) – 1.5 x supernatant cholesterol (mmol/L).

High density lipoprotein (HDL)

Principle

LDL and VLDL are precipitated from serum by the action of a polysaccharide in the presence of divalent cations. Then, HDL present in the supernatant is determined.

Procedure

The precipitant solution, 0.1 ml was added to 0.3 ml of the serum sample and mixed thoroughly and allowed to stand for 15 min. This was centrifuged at 2,000 xg for 15 min. The cholesterol concentration in the supernatant was determined. Determination of the concentration of the serum total HDL as described by Kameswara et al. (1999) was used.

Triacylglycerol


Clinical significance

Triacylglycerols measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g diabetes mellitus, nephrosis and liver obstruction.

Principle

The triacylglycerols are determined after enzymatic hydrolysis with

lipases. The indicator is a quinoneimine formed from hydrogen peroxide, 4-aminophenazone and 4-chlorophenol under the catalytic influence of peroxidase.



Method

A quantity of the sample (0.1 ml) was pipetted into a clean labeled tube and 1.0 ml of trichloroacetic acid (TCA) was added to it, mixed and then centrifuged at 250 rpm for 10 min. The supernatant was decanted and reserved for use.

The mixtures were allowed to stand for 20 min at 25°C and the absorbance of the sample and standards read against the blank was taken at 540 nm.

Calculation

The concentration of triacylglycerol in serum was calculated as follows:

$$\frac{\text{Absorbance of sample}}{\text{Absorbance of standard}} \times \text{Standard concentration (mmol/l)} = \text{mmol/l}$$

Statistical analysis

Statistical analysis was carried out using Statistical Package for Social Sciences (SPSS) version 16.0. One way analysis of variance was adopted for comparison, and the results were subject to post hoc test using least square deviation (LSD). The data were expressed as mean \pm standard deviation. $P < 0.05$ was considered significant.

RESULTS

The effects of tyrosine supplemented diet on the total cholesterol levels of rats

There was significant increases ($p < 0.05$) in total cholesterol of week 2 of rats fed with 0.8 g/100 g of tyrosine as compared to that fed with 1.0 g/100 g and 1.2 g/100 g of the tyrosine supplemented diet. There was also a significant increase ($p < 0.05$) of rats fed with 1.0 g/100 g and 1.2 g/100 g of tyrosine meal diet. The same trend was also observed in week 3 (Figure 1). This increases were not dose dependent.

The effects of tyrosine supplemented diet on the HDL levels of rats

The bar shows the result of the HDL of rats fed with tyrosine supplemented diet. There was significant




Figure 1. Bar chart representing the effects of tyrosine meal diet on the total cholesterol levels of rats. Values indicates mean \pm SEM (n=4). Significance is at $p < 0.05$.




Figure 2. Bar chart representing the effects of tyrosine meal diet on the HDL levels of rats. Values indicates mean \pm SEM (n=4). Significance is at $p < 0.05$.

decrease ($p < 0.05$) in HDL when rats fed with 0.8 g/100 g of tyrosine meal diet was compared to the normal control of week 2. There was no significant difference ($p > 0.05$) in weeks 2 and 3 rats fed with 0.08 g/100 g tyrosine supplemented diet when compared with 1.0 g/100 g and 1.2 g/100 g meal diet (Figure 2).

tyrosine supplemented diet. There was significant increases ($p < 0.05$) in week 2 of the LDL of rats fed with 1.0 g/100 g and 1.2 g/100 g tyrosine supplemented diet as compared to the control. Significant increases ($p < 0.05$) was also observed in of rats fed with 0.8 g/100 g, 1.0 g/100 g and 1.2 g/100 g tyrosine meal diet at week 3 when compared with the control (Figure 3).

The effects of tyrosine supplemented diet on the LDL levels of rats

DISCUSSION

The bar shows the result of the LDL of rats fed with

The effects of dietary tyrosine on serum lipid profile were




Figure 3. Bar chart representing the effects of tyrosine meal diet on the LDL levels of rats. Values indicate mean \pm SEM (n=4). Significance is at $p < 0.05$.

analyzed in this study. The animals fed with 0.8 g/100 g, 1.0 g/100 g and 1.2 g/100 g of tyrosine supplemented diet showed significant increases ($p < 0.05$) in total cholesterol of week 2 of rats as compared to that fed with 1.0 g/100 g and 1.2 g/100 g of the tyrosine supplemented diet. There was also a significant increase ($p < 0.05$) in rats fed with 1.0 g/100 g and 1.2 g/100 g of tyrosine meal diet. The same trend was also observed in week 3 as reflected in this work. Increased total cholesterol concentration when fed with high tyrosine meal diet allows one to investigate the effects of treatment in dietary protein source (tyrosine), on this fraction. These results followed a similar trend in the rise of total cholesterol levels using other sources of proteins (soy protein and casein) as reported by other researchers (Forsythe et al., 1980; Nagata et al., 1982c; Van der Meer, 1983; Van der Meer et al., 1985). Our results showed a significant elevation of cholesterol in the serum of experimental rats as compared to the control. Cholesterol is synthesized in the liver; this result demonstrates the ability of tyrosine to influence liver metabolism towards increased synthesis of lipids. The high levels of cholesterol may be due to a number of factors such as the increased availability of fatty acids for esterification (Bopama et al., 1997), reduced catabolism of LDL, inhibition of tissues, activation of acetyl-CoA carboxylase (McCarthy, 2001) and production of triglyceride precursors such as acetyl-CoA and glycerol phosphate (Fatiha et al., 2014). The elevation of cholesterol in the liver might suggest that the

acid synthase, glucose 6-phosphate dehydrogenase and HMG-CoA reductase (Vega et al., 2003) which are required for cholesterol synthesis.

Atherosclerosis is characterized by liver disease without alcohol which is manifested by the significant lipid deposition in hepatocytes of liver parenchyma as a single macro-vesicular steatohepatitis and can develop fibrosis in cirrhosis which is increasingly recognized as an important cause of mortality (Angulo and Lindor, 2002).

There was a significant decrease ($p < 0.05$) in HDL in rats fed with 0.8 g/100 g of tyrosine supplemented diet as compared to the control of week 2. There was no significant ($p > 0.05$) increase in weeks 2 and 3 of rats fed with 0.8 g/100 g tyrosine meal diet when compared with 1.0 g/100 g and 1.2 g/100 g supplemented diet. In the present study, in the rats fed higher doses of tyrosine serum HDL continued to have a higher amount as in the case of total cholesterol as compared to the normal rats fed supplemented tyrosine diet during the 21-day study. These observations were in contrast with the work of Nagaoka et al. (1990a) who reported that prolonged administrations of dietary tyrosine caused hypercholesterolemia in male Wistar rats. There were significant increases ($p < 0.05$) in week 2 of the LDL of rats fed with 1.0 g/100 g and 1.2 g/100 g tyrosine supplemented diet when compared with the control. Significant increases ($p < 0.05$) were also observed in week 3 of rats fed with 0.8 g/100 g, 1.0 g/100 g and 1.2 g/100 g tyrosine supplemented diet when compared with the

dietary tyrosine supplement contain ingredients capable of enhancing the activities of hepatic lipogenic and cholesterologenic enzymes, such as malic enzyme, fatty

control. Excess dietary tyrosine or certain xenobiotics increases HDL and LDL cholesterol (Kato and Yoshida, 1981; Nagaoka et al., 1985b). In the hypercholesterolemia

seen following feeding of cholesterol, serum LDL cholesterol was relatively increased (Quazi et al., 1983). It seems likely that cholesterol metabolism is quite different in rats fed higher doses of tyrosine from those of rats fed cholesterol-containing diet. Many investigators have already suggested that the hypercholesterolemia seen in feeding dietary tyrosine (Qureshi et al., 1978; Solomon and Geison, 1978) was mediated by enhancement of of cholesterol synthesis in the liver.

CONCLUSION

The alterations in the serum cholesterol reveals that long term feeding of dietary tyrosine may escalate cholesterol accumulation in the adipose tissues.

Conflict of Interests

The author(s) have not declared any conflict of interests.

REFERENCES

- Abell LL, Levey BB, Brodie BB, Kendall FE (1952). Extraction of Cholesterol. *J. Biol. Chem.* 195(1):357-366.
- Angulo P, Lindor KD (2002). Nonalcoholic fatty liver disease. *N. Engl. J. Med.* 346:1221-1231.
- Bopama KH, Kannan J, Gadgil S, Balaraman ER. (1997). Antidiabetic and antihyperlipidemic effects of neem seed kernel powder on alloxan diabetic rabbits. *Indian J. Pharmacol.* 29:162-167.
- Deijen JB, Orlebeke JF (1994). Effect of tyrosine on cognitive function and blood pressure under stress. *Brain Res. Bull.* 33(3): 319–323.
- Deijen JB, Wientjes CJ, Vullinghs HF, Cloin PA, Langefeld JJ (1999). Tyrosine improves cognitive performance and reduces blood pressure in cadets after one week of a combat training course. *Brain Res. Bull.* 48(2):203-209.
- Elwes RD, Crewes H, Chestman T (1989). Treatment of narcolepsy with L-tyrosine: double-blind placebo-controlled trial. *Lancet* 2:1067-1069. Fatiha O, Noreddine G, Mohamed E, Hakima B, El-Mustapha D, Souliman A (2014). Hypolipidemic and anti-atherogenic effect of methanol extract of Fennel (*Foeniculum vulgare*) in hypercholesterolemic mice. *Int. J. Sci. Knowl.* 3(1):42-52.
- Fernstrom JD (2000). Can nutrient supplements modify brain function? *Am. J. Clin. Nutr.* 71(6 Suppl):1669-1675.
- Forsythe WA, Miller ER, Hill GM, Romsos DR, Simpson RC (1980). Effects of dietary protein and fat sources on plasma cholesterol parameters, LCAT activity and amino acid levels and on tissue lipid content of growing pigs. *J. Nutr.* 110:2467-2479.
- Gordon DJ, Probstfield JL, Garrison RJ, Neaton JD, Castell, WP, Knoke JD, Jacobs DR, Bangdiwala S, Tyroler HA (1989). High-density lipoprotein cholesterol and cardiovascular disease. Four prospective American studies. *Circulation* 79(1):8-15.
- Kameswara RB, Kesavulu MM, Giri CH (1999). Anti-diabetic and hypolipidemic effects of *Momordica cymbalaria* Hook fruit powder in alloxan-induced diabetic rats. *J. Ethnopharmacol.* 67:103 -109.
- Kato N, Yashida A (1981). Effects of various dietary xenobiotics on serum total cholesterol and high density lipoprotein cholesterol in rats. *Nutr. Rep. Int.* 23:825-831.
- Lewis GF, Rader DJ, (2005). New insight into the regulation of HDL metabolism and reverse cholesterol transport. *Circ. Res.* 96(12): 1221-1232.
- McCarthy MF (2001). Inhibition of acetyl-CoA carboxylase by cystamine may mediate the hypotriglyceridemic activity of pantethine. *Med. Hypotheses* 56(3):314-317.
- Moller SE, Maach-Moller B, Olesen M, Madsen P, Fjalland B (1995). Tyrosine metabolism in users of oral contraceptives. *Life Sci.* 56:687- 695.
- Nagaoka S, Aoyama Y, Yoshida A (1985b). Effect of tyrosine and some other amino acids on serum level of cholesterol in rats. *Nutr. Rep. Int.* 31:1137-1148.
- Nagaoka S, Miyazaki H, Oda H, Aoyama Y, Yoshida A (1990a). effects of excess dietary tyrosine on cholesterol, bile acid metabolism and mixed-function oxidase system in rats. *J. Nutr.* 120(10):1134-1139.
- Nagata Y, Tanaka K, Sugano M (1982c). Further studies on the hypercholesterolemic effect of soy protein product in rats. *Br. J. Nutr.* 45:855-862.
- Olson RE (1998). Discovery of the lipoproteins, their role in fat transport and their significance as risk factor. *J. Nutr.* 128(2):439-443.
- Quazi S, Yokogoshi H, Yoshida A (1983). Effect of dietary fiber on hypercholesterolemia induced by dietary PCB or cholesterol in rats. *J. Nutr.* 113:1109-1118.
- Qureshi AA, Solomon JK, Eichelman B (1978). L-Histidine induced fascilitation of cholesterol biosynthesis in rats. *Proc. Soc. Exp. Biol. Med.* 159:57-70.
- Salter CA (1989). Dietary tyrosine as an aid to stress resistance among troops. *Mil. Med.* 154(3):144-146.
- Shurtleff D, Thomas JR, Schrot J, Kowalski K, Harford R (1994). Tyrosine reverses a cold-induced working memory deficit in humans. *Pharmacol. Biochem. Behav.* 47:935-941.
- Sole MJ, Benedict CR, Myers MG, Leenen FH, Anderson GH (1985). Chronic dietary tyrosine supplements do not affect mild essential hypertension. *Hypertension* 7(4):593-596.
- Solomon JK, Geison RL (1978). Histidine-induced hypercholesterolemia: characteristics of cholesterol biosynthesis in rat livers. *Proc. Soc. Exp. Biol. Med.* 159:44-47.
- Thomas JR, Lockwood PA, Singh A, Deuster PA (1999). Tyrosine improves working memory in a multitasking environment. *Pharmacol. Biochem. Behav.* 64(3):495-500.
- Tumilty L, Davison G, Beckmann M, Thatcher R (2011).

- Oral tyrosine supplementation improves exercise capacity in the heat. *Eur. J. Appl. Physiol.* 111(12):2941-2951.
- Van der Meer R (1983). Is the hypercholesterolemic effect of dietary casein related to its phosphorylation state? *Atherosclerosis* 49:339- 341.
- Van der Meer R, De Vries H, West CE, Dewaard H (1985). Casein- induced hypercholesterolaemic in rabbits in calcium-dependent. *Atherosclerosis* 56:139-147.
- Vega GL, Weiner MF, Lipton AM. (2003). Reduction in levels of 24S- hydroxycholesterol by statin treatment in patients with Alzheimer disease. *Arch. Neurol.* 60:510-515.